efit. In this way its external use is sometimes serviceable.

In old syphilitic affections, it is occasionally useful. It has been recommended in hooping cough, but it is not a perfectly safe medicine for children, owing to the difficulty of ascertaining when its constitutional effects take place in them.—I am informed on the best medical authority, that it is of great use in some cases of hemicrania, which are not regularly intermittent.

The most common form of preserving the Hemlock for use, as well as the most convenient for its exhibition, is that of the inspissated juice or extract. It is well known however, that the extracts kept in our shops differ materially in their strength, so that in beginning from a new parcel, the physician can seldom predict the degree of operation of his first doses. In some instances very great quantities have been taken without the least effect. The extract is apt to prove inert when the plant is gathered too young, when the evaporation is conducted with too much heat, when a decoction of the dried plant has been evaporated instead of the fresh leaves, and lastly when the extract itself has become old. To give the extract its due strength, the plant should be collected at full maturity, while in flower, or in fruit provided it remains green, and

the juice or the decoction should be evaporated at the heat of boiling salt water. The stock should be renewed every year. A suitable dose for commencement is from one to five grains. This may be increased at every time of taking it, until its constitutional effects are felt. In beginning the use of a new parcel, more caution is requisite at first, than after its strength has been tested.

The Æthusa cynapium, an umbellate plant very common in Boston, has sometimes been mistaken for Hemlock, which it considerably resembles. It is a smaller plant, with its stalk not spotted. It differs also in having no general involucre, while its partial one is very long.

BOTANICAL REFERENCES.

Conium Maculatum, Linnæus. *Sp. Pl.*—Woodville, *t.* 22.—Curtis, *Fl. Lond.* i. 17.—Smith, *Engl. Bot. t.* 1191.—Pursh, i. 195.—Cicuta vulgaris, Morison, *Umb. t.* 6.—Parkinson, *Theatr.* 933.—Cicuta major, Lamarck, *Encyclopedie Methodique.*—Cicutaria vulgaris, Clusius, *Hist.* 200.

MEDICAL REFERENCES.

Murray, *Apparatus medicaminum*, i. 322.—Cullen, *Mat. Med.* ii. 263.—Fothergill, *Med. Obs.* iii. 400.—Hunter *on the venereal*, 108, 175, 199, 247, &c.—Home, *Annals Med.* iii. 66.—Butter, *Med. Comment.* i. 373.—Fisher, *Mem. Mas. Soc.* i.—Jackson, *N. Engl. Journ.* ii. 105.—Guersent. *Dict. Sciences Medicales*, v. 208.—Orfila, *Toxicologie*, iii. 279, &c. &c. &c.